

Text in Red, if correct, leave it in black.
Inform the author that it will be in the header.
Correct the references.

Article - Biological and Applied Sciences

Survival of Amputated Striped Corvina *Cynoscion reticulatus* (Pisces: Sciaenidae) off the Southeast Coast of the Gulf of California

Juan Antonio Maldonado Coyac¹

<https://orcid.org/0000-0001-6535-0405>

Juan Ángel Payán Alcacio²

<https://orcid.org/0000-0003-2402-4528>

Maria de los Angeles Maldonado Amparo¹

<https://orcid.org/0000-0003-3005-9348>

Marcelino Ruiz Domínguez^{1*}

<https://orcid.org/0000-0001-9632-8008>

¹Universidad Autónoma de Sinaloa, Facultad de Ciencias del Mar, Mazatlán, Sinaloa, México; ²Instituto Politécnico Nacional, Centro Interdisciplinario de Ciencias Marinas, La Paz, Baja California Sur, México.

Editor-in-Chief: Paulo Vitor Farago
Associate Editor: Paulo Vitor Farago

Received: 28-Nov-2022; Accepted: 15-Sep-2023

*Correspondence: marcelinoruizdom@uas.edu.mx.com; Tel.: 6121676130 (M.R.D.).

HIGHLIGHTS

- Striped corvina were able to swim and compete for resources despite amputation.
- Results could be useful in trophic and swimming modeling, and fitness survival.

Abstract: Records of amputated teleost fish in the aquatic environment are becoming more frequent; however, it is difficult to find the causes of amputation when there is little evidence. For this reason, this study focused on describing the damage and causes of the amputations of two striped corvina *Cynoscion reticulatus* specimens collected from landing sites of a small-scale fishery in northwestern Mexico, and determining if the damaged organisms could survive in the environment. Specimens were sexed and whole-body radiographs of amputated specimens and of a normal specimen were taken. Morphometric measurements (TL, SL, MDF, BD, HL, and TW) of the amputees and of 54 normal specimens were recorded. The TL of amputees was estimated and compared to that of normal specimens; stomach contents were also analyzed. The first organism was male, it was missing 1 ½ vertebra and had an estimated TL of 36.64 cm; the second was a female, it was missing 3 ½ vertebrae, with an estimated TL of 38.59 cm; both had bite marks, skin regeneration, and scales in the affected area. Stomach analysis showed Engraulidae fish of the *Anchoa* genus in stomach contents. There were no significant differences in the measurements of the two amputated and 54 normal organisms ($p>0.05$). Based on this evidence, it was determined that amputated fish could survive, escape from predators, feed, and obtain energy to heal wounds; the amputations were

attributed to the bottlenose dolphin *Tursiops truncatus*, a potential predator of croakers in the study area and other regions.

Keywords: injuries; tailless; healing; predator bite; small-scale fishery.

INTRODUCTION

Fish suffer various anatomical injuries in the aquatic environment (e.g., loss of scales, spines, rays, fins, and vertebrae) [1, 2] due to pathogenic organisms such as bacteria, fungi [3], and parasites [4, 5, 6]; bites from fish and marine mammals [7, 8]; and fishing lines or nets [9], and polluting plastics such as six-pack rings and remains of fishing gear [10]. All of these conditions, plus interspecific and intraspecific competition and environmental conditions, result in constant stress on organisms.

The most commonly reported injury in recent studies on free-living marine fish has been fin loss, especially the caudal fin and peduncle. However, it has been reported that fish have developed physiological abilities that allow them to heal their wounds in a short time. For example, there are reports of two specimens of striped piggy *Pomadasys stridens* (Forsskal, 1775) in the Persian Gulf [7] and two croakers *Micropogonias undulatus* (Linnaeus, 1766) from Biloxi Bay and the Mississippi Sound [1] presenting amputation of the fin and caudal peduncle; however, the affected area presented scarring and was covered by scales. Additionally, it has been reported that upon suffering amputation of the caudal fin, morid cod *Physiculus cyanostrophus* [11] from the eastern tropical Atlantic [12] and zebrafish *Danio rerio* (Hamilton-Buchanan, 1822) [13] regenerated the fin completely after 2 to 3 weeks. Although fish can heal their wounds, the loss of a caudal fin makes feeding difficult, it decreases body condition, the efficiency of escape from predators, and affects other daily activities [14; 15], as this appendage provides the fish with thrust and direction during swimming [16]. Many of the reports on amputations in wild fish highlight the difficulty of defining the origin of these injuries when there is little evidence, and it is recommended to address such explanations based on the biological knowledge of the species, environmental conditions, and the anthropogenic activity in the area where the records were made.

The present study documents for the first time the presence of two striped corvina *Cynoscion reticulatus* (Günther, 1864) specimens lacking a caudal fin and caudal peduncle, found by fishers in southern Sinaloa. This croaker species is distributed from Baja California Sur, Gulf of California, to Colombia [17], and it is an important socioeconomic resource along its distribution area [18, 19, 20]. For more than 20 years, *Cynoscion reticulatus* has been part of a Mexican artisanal fishery that uses 3" to 6" mesh size nets of variable length (100 to 500 m); it is also captured as bycatch by shrimp boats using trawls, gillnets, and hand lines [21]. Regarding its ecology, it is a fourth-level predator (4.0 ± 0.24), a carnivore, and its preferred diet includes fish, shrimp, and cephalopods [22]. It is preyed upon by large fish such as groupers, sharks, and rays [23], as well as some marine mammals, such as the sea lion *Zalophus californianus* (Lesson, 1828) [24, 25] and the vaquita *Phocoena sinus* (Norris & McFarland, 1958) [8] in the Gulf of California and to the southeast of this region. Therefore, the objective of this work was to describe the injuries, body condition, and most probable cause of the amputations. The following questions were raised: 1.-Which of all the factors described as causing amputations in fish could be the one that affected the striped corvinas? 2.- Could the amputated croakers have survived and developed correctly in their environment? To answer these questions, the following hypotheses were raised: 1. The main cause of damage to the fin and caudal peduncle of striped corvinas was due to a bite from one of their predators; 2. Amputated striped corvinas can survive in the environment as long as they can feed and maintain good physical condition.

MATERIAL AND METHODS

Two sampling trips were undertaken in August and October 2021; specimens were obtained from small-scale fishery landings in southern Sinaloa. This fishery targets fish in the area comprising from Punta Prieta (23°36'N, 106°54'W) to Chametla (22°45'N, 106°02'W), using >300 m long gillnets with 3" and 3 ½" mesh size, at depths below 50 m. Two tailless specimens and 54 normal (non-amputated) specimens of the striped corvina *Cynoscion reticulatus* (n= 25 August 2021 and n= 29 October 2021) were obtained (Figure 1a). These fishers have permits for commercial fishing of marine finfish, authorized by the National Commission for Fisheries and Aquaculture through the Secretary of Agriculture and Rural Development of the Government of Mexico; since the fish were already dead when they were collected, no permit was required to handle them.




Figure 1. Photographic comparison of a tailless (a) male and (b) female versus a tailed (c) female *Cynoscion reticulatus* from the southeast Gulf of California.

The total length (TL), standard length (SL), length from the mouth to the end of the second dorsal fin (MDF), body depth (BD), head length (HL), and total weight (TW) were obtained for all specimens. Detailed photographs of the amputation area of both specimens were taken, (Figure 2) and X-ray images (Figure 3) of the two tailless and one tailed fish to describe possible injuries to the bone system. The specimens were dissected to determine sex by gonadal visualization and the stomach was removed to verify if the amputated fish could feed despite their condition. The taxonomic analysis of the target species and of prey was based on the keys by [17, 26]. The Total length of the two amputated organisms was estimated using, the average percentage represented by the MDF with respect to the TL of the 54 non-amputated organisms, and the MDF was extrapolated to the TL using the percentage mean result as a factor:

$$Factor = \left(\frac{\frac{TL_1}{MDF_1} + \frac{TL_2}{MDF_2} + \frac{TL_3}{MDF_3} + \frac{TL_4}{MDF_4} + \dots + \frac{TL_n}{MDF_n}}{n} \right) * 100$$

Where $n=54$ non-amputated organisms. After calculating the factor, the following arithmetic operation was performed.

$$LT_{Amputated} = MDF_{Amputated} \times Factor$$

Finally, the measurements (TL, MDF, BD, HL, and TW) of the two tailless organisms and of the 54 normal croakers were compared (Table 1) using a *t*-student test for independent samples [27].

RESULTS

The two amputated organisms were dissected, and the external and internal morphologies were analyzed. Based on the macroscopic observation of the gonads, we determined that the first tailless specimen found in August 2021 was a sexually immature male presenting signs of amputation of the caudal fin. We observed that the dermal tissue of the injured area showed recent scarring, and the absence of scales was evident; instead of scales, there was increasing pigmentation of the dermis, although the vertebral bone tissue could be seen through the skin (Figure 2a). The second specimen lacking a caudal fin was found in October 2021 (Figure 1b); it was identified as a sexually mature female (the ovaries presented secondary-growth oocytes). This specimen presented advanced scarring of the amputated area, slight skin pigmentation, and like in the first specimen, the vertebral tissue could be observed through the muscular tissue (Figure 2b).




Figure 2. Amputated region of *Cynoscion reticulatus* (a) male and (b) female specimens from the entrance to the Gulf of California. 1. Vertebral bone tissue; 2. Pigmentation of scar tissue; 3. Evidence of bite or other tissue damage.

The analysis of the X-ray image showed that the first specimen (August 2021) had 13 precaudal vertebrae (PrCv) and 10 ½ caudal vertebrae (Cv), for a total of 23 ½ vertebrae (Figure 3a). We could also observe bone damage, recovery of scale cover in the amputated area, and healing in spines V, VI, X, and XI, as well as in rays 1-8 of the dorsal fin, as shown by the white dotted circles 1 and 2 in Figure 3a. This image also allowed us to identify the presence of food in the stomach; the dissection showed a pair of small sagitta otoliths and two fish from the family Engraulidae. The entire contents (otoliths and fish) were identified as anchovies from the genus *Anchoa*. The second specimen (October 2021) had a total of 13 PrCv and 8 ½ Cv, for a total of 21 ½ vertebrae (Figure 3b). Regarding stomach contents, only two pairs of small sagitta otoliths belonging to fish from the family Engraulidae, genus *Anchoa*, could be found. The estimated factor for the extrapolation of the TL of the two amputated striped corvinas was 1.308 ± 0.024 ; therefore, the estimated TL for the first striped corvina was 36.64 cm and 38.59 cm for the second.

The third analyzed striped corvina (not amputated) was a sexually mature female measuring 36.56 cm TL (Figure 1c). This specimen was used as reference for the bone structure of a specimen not lacking appendages. According to the X-ray image, it presented 13 PrCv and 12 Cv, for a total of 25 vertebrae (Figure 3c). The stomach analysis showed stomach contents (one fish with 18 vertebrae at an advanced state of digestion that could not be identified), which can be observed in Figure 3a, shown by a dotted circle in the abdominal region.




Figure 3. X-ray comparison of normal and tailless *Cynoscion reticulatus* specimens from the entrance to the Gulf of California. a) Tailless female (white arrow) with bone damage to the spines and rays (dotted circles 1 and 2) and presence of food in the stomach (dotted circle 3); b) Tailless male (white arrow); c) Normal female with presence of food in the stomach (dotted circles).

Finally, the comparison of measurements (TL, MDF, BD, HL, and TW) of the two tailless organisms and the 54 normal croakers did not show significant differences ($P > 0.05$; Table 1).

Table 1. Morphology of normal and amputated organisms of striped corvina *Cynoscion reticulatus*.

Measurements (cm)	1 st tailless male	2 nd tailless female	Average size of normal fish
TL	36.64	38.59	37.18 ± 3.54
SL			32.5 ± 3.31
MDF	28.01	29.51	28.37 ± 2.95
BD	7.23	7	7.79 ± 0.81
HL	8.89	9.7	9.08 ± 0.92
TW	215	321	488.61 ± 138.85

DISCUSSION

The characteristics of the two striped corvinas *Cynoscion reticulatus* that were lacking a tail or caudal peduncle (e.g., mid vertebrae, lack of complete vertebrae, scarring, pigmentation, damage to spines and dorsal rays, and bite features in the damaged area) indicated that the main cause of injury was due to a bite from one of their predators (e.g. grouper, shark, ray, sea lion, vaquita, or dolphin) [8, 24, 25], the distribution and hunting strategy of these predators were reviewed to found the most probable one. The vaquita was ruled out since its distribution area is limited to the upper Gulf of California [28], in addition, as the wounds of the striped corvina were recent, and their swimming ability impaired, it would not be able to move that far (Southeast of the Gulf of California). Groupers and rays were also discarded, it is unlikely that they can cut the appendices of the medium fishes due to the type of teeth they have (small canines in groupers and dental plates in rays) [29, 30], in addition their feeding strategy is based on suction, which would otherwise favor the capturing of the entire prey [31, 32]. Although there are various hunting strategies within elasmobranchs, from the possible ones related to depth, some specialize in suction in the benthic zone, while most epibenthic and pelagic elasmobranchs with generalist habits use the lunge, suck, and bite strategy to catch prey [33]. Considering the type of dentition of the sharks that inhabit the Southeastern region of the Gulf of California (*Sphyrna spp.*, *Rhizoprionodon longurio*, *Carcharhinus spp.*, *Nasolamia velox*, *Galeocerdo cuvier*) [17], the attack strategy and biting pattern they use to feed, we deduce that this group is among the possible causes of amputations. Another viable species would be the sea lion (*Zalophus californianus*) and the bottlenose dolphin (*Tursiops truncatus*), both use ramming (impulse towards the prey with the mouth semi-closed and open) as their feeding strategy, as well as cooperative hunting through grouping for more efficiently feeding [34, 35, 36], being able to bite and tear prey. Therefore, this would respond to the first hypothesis.

The evidence found in the two amputated croakers indicated that the bite was recent and that the specimens were part of the same population, because in addition to the fact that the wounds were healed, their condition was similar to that of normal croakers. This was confirmed by the fact that no significant differences were found when comparing body dimensions (TL, MDF, BD, HL, and TW), and was also corroborated by the diet analysis, as both amputated and normal croakers presented food remains in the stomach. These results were decisive to accept the second hypothesis. Responding to the second hypothesis, this would demonstrated that although the croakers had a swimming disability, they could compete for the same resources in the environment where they were found, which could be understood as a benefit of gregarious intraspecific relationships (e.g., fish schools).

Fish anomalies have been reported sporadically in the past. For the study area, in the Urias Estuary in Mazatlán, Sinaloa (an area adjacent to the study area), the absence of pelvic fins in a yellow snapper *Lutjanus argentiventris* (Peters, 1869) was reported, highlighting the fact that the lack of fins was due to congenital causes and not predation, fishing gear, or environmental factors, because when compared with a normal fish from the same species, it had the same physical condition and anatomical organization, except for the absence of pelvic fins [37]. Regarding reports of injuries in croakers, there is a study by [1] on three Atlantic croakers *Micropogonias undulatus* caught with trawl nets in Biloxi Bay and the Mississippi Sound, which presented amputation of the caudal fin and peduncle, with scarring and covering with scales of the affected area (similar to what was found in the present study). This damage was attributed to bites from predators, as they were not the only fish found with bites. The authors also found a tailless Atlantic spade fish *Chaetodipterus faber* (Broussonet, 1782); two Gulf menhaden *Brevoortia patronus* (Goode, 1878), the first with a halfmoon bite on the abdomen and the other with a lesion on the back, both with scarring and absence of scales in the affected area; and a tailless female lined seahorse *Hippocampus erectus* (Perry, 1810) with scarring of the wound. Recently, the Texas Parks and Wildlife reported a tailless spotted seatrout *Cynoscion*

nebulosus (Cuvier, 1830) lacking a caudal peduncle, but presenting a developed stump covered with scales; however, the causes of the injury were not described (<https://tpwd.texas.gov/fishboat/fish/didyouknow/coastal/amazingfish.phtml>).

These conditions have also been identified in other species; for example, in the Bay of Bengal in the Indian Ocean, [38] an orange-spotted grouper *Epinephelus coioides* (Hamilton, 1822) presenting a stump in the caudal region and a Bengal tongue sole *Cynoglossus cynoglossus* with a small caudal fin in regeneration were recorded; both fishes showed signs of having suffered recent damage. However, causes were attributed to the environment, as there are several reports of fish with abnormalities in that study area.

Another specific condition known as "fin rot" or "fin erosion" disease in fish, associated with degraded coastal or estuarine environments, has been reported in the weakfish *Cynoscion regalis* and summer flounder *Paralichthys dentatus* from the Atlantic Ocean [39]. *Vibrio* sp. has been described as a possible determinant of erosions in the anal, caudal, and pelvic fins of weak fish; as well as erosions in large areas of the edges of the fins of the summer flounder.

Finally, if potential predators are considered, among the ones found in the region, sea lions, sharks, and bottlenose dolphins, the latter might be the most likely candidate for the amputations found in the striped corvina, at least, for the southern Gulf of California, as it has been recorded as one of the top predators of corvina for other areas in the Pacific Northwest [40], Southern Africa [41], the Southeastern United States [42], and the Pacific coast of South America [43]. Adding to this hypothesis, it has been reported that in the Gulf of Mexico (Laguna Tamiahua, Veracruz), the bottlenose dolphin developed a particular hunting strategy, it only bites the posterior body area of catfish (*Ariopsis felis* and *Bagre marinus*) to avoid choking on the dorsal and pectoral fin spines [44]. The results of this work can be useful in studies focused on the hunting strategies of the main predators of striped corvina or similar members of the scianidae family, ecosystem trophic modelling, swimming capacity modelling, healing capacity and survival of this and other corvinas species.

Funding: This research received no external funding.

Acknowledgments: We are grateful for Project PROFAPI2022 "PRO_A7_012" granted by the Universidad Autónoma de Sinaloa.

Conflicts of Interest: The authors declare no conflict of interest.

- All citations must appear in the list of references, and vice-versa;

- References - Format correctly (abbreviation of authors, journal title, etc.).

- In the references, if there is a link, are accessing it correctly?

REFERENCES – abbreviate Journals' titles, the title of the article if it is in another language, translate it into English and leave it in brackets.

1. Gunter G, Ward JW. Some fishes that survive extreme injuries, and some aspects of tenacity of life. *Copeia*. 1961;4:456-62. Available from: <http://www.jstor.org/stable/1439589>
2. Jawad LA, Ibrahim M, Waryani B. Incidences of caudal fin malformation in fishes from Jubail City, Saudi Arabia, Arabian Gulf. *Fish Aquat Life*. 2018;26:65-71. <https://doi.org/10.2478/aopf-2018-0008>
3. Austin B, Austin DA. Bacterial fish pathogens: disease of farmed and wild fish. Vol. 26. Chichester: Springer; 2007.
4. Çelik S, Korun J, Gökoğlu M. First occurrences of *Nerocila bivittata* on Dusky Grouper (*Epinephelus marginatus*) and Mottled Grouper (*Mycteroperca rubra*). *J Hellenic Vet Med Soc*, 2020;71(3):2309-2314. <https://doi.org/10.12681/jhvms.25077>

5. Aguilar-Perera A. Checklist of Parasitic Isopods (Crustacea: Isopoda) Infesting Marine Decapod and Fishes Off Mexico's Coasts. *Thalassas*. 2021. <https://doi.org/10.1007/s41208-021-00336-x>
6. Williams EH, Bunkley-Williams L. Life cycle and life history strategies of parasitic crustacea. In: Smit NJ, Bruce NI, Hadfield KA, editors. *Parasitic Crustacea: State of Knowledge and Future Trends*. Switzerland: Springer Nature; 2019. p. 179-266. https://doi.org/10.1007/978-3-030-17385-2_5
7. Alavi-Yeganeh MS, Razavi S, Egan JP. Taillessness and skeletal deformity in striped piggy *Pomadasys stridens* (Osteichthyes: Haemulidae) from the Persian Gulf. *Dis Aquat Organ*. 2019;132:209-13.
8. Pérez-Cortés MH, Silber GK, Ramírez BV. Contribución al conocimiento de la alimentación de la vaquita, *Phocoena sinus*. *Cienc Pesq*. 1996;13:66-77.
9. Richardson K, Asmutis-Silvia R, Drinkwin J, Gilardi K, Giskes I, Jones G, et al. Building evidence around ghost gear: Global trends and analysis for sustainable solutions at scale. *Mar Poll Bull*. 2019;138:222-9. <https://doi.org/10.1016/j.marpolbul.2018.11.031>
10. Andrady AL. Plastics in the Anthropocene. In: Andrady AL, editor. *Plastics and the Ocean: Origin, Characterization, Fate, and Impacts*. Nueva Jersey: Wiley; 2022. p. 1-42.
11. Anderson ME, Tweddle D. A new species of *Physiculus* (Teleostei: Moridae) from the southeastern Atlantic. *Arch Fish Mar Res*. 2002;50(1):17-22.
12. Triay-Portella R, González JA, Pajuelo JG. Caudal region regeneration in a natural population of the morid fish *Physiculus cyanostrophus* in the tropical eastern Atlantic ocean. *Deep-Sea Res I: Oceanogr Res Pap*. 2019;150:103062. <https://doi.org/10.1016/j.dsr.2019.06.008>
13. Sehring IM, Weidinger G. Recent advancements in understanding fin regeneration in zebrafish. *Wiley Interdiscip Rev Dev Biol*. 2019;e367. <https://doi.org/10.1002/wdev.367>
14. Böckelmann PK, Ochandio BS, Bechara IJ. Histological study of the dynamics in epidermis regeneration of the carp tail fin (*Cyprinus carpio*, Linnaeus, 1758). *Braz J Biol*. 2010;70(1):217-23. <http://dx.doi.org/10.1590/s1519-69842010000100030>
15. Fu C, Cao ZD, Fu SJ. The effects of caudal fin loss and regeneration on the swimming performance of three cyprinid fish species with different swimming capacities. *J Exp Biol*. 2013;216(16):3164-3174. <https://doi.org/10.1242/jeb.084244>
16. Flammang BE, Lauder GV. Speed-dependent intrinsic caudal fin muscle recruitment during steady swimming in bluegill sunfish, *Lepomis macrochirus*. *J Exp Biol*. 2008;211(4):587-598. <https://doi.org/10.1242/jeb.012096>
17. Robertson DR, Allen GR. [Tropical Eastern Pacific Coastal Fishes: Online Information System. Version 2.0 Smithsonian Tropical Research Institute, Balboa, Republic of Panama. cited 2022 Jul 14]. Available from: <https://biogeodb.stri.si.edu/sftep/es/thefishes/species/1526>

18. Saucedo-Barrón CJ, Ramírez-Rodríguez M. [Commercially Important Fish in the Southern Region of Sinaloa State, Mexico (Artisanal Fishing)]. Invest Mar CICIMAR. 1994;9(1):51-54.
19. Musso-Solari MB. [Reproductive Cycle of the Striped Corvina *Cynoscion reticulatus* (Günther, 1864), and the Croaker *Micropogonias ectenes* (Jordan and Gilbert, 1882) (Pisces: Sciaenidae) from the Coast of Sinaloa] [Dissertation]. Instituto de Ciencias del Mar y Limnología: Universidad Nacional Autónoma de México; 2011.
20. Ortiz JR, Pacay-Barahona AJ, Polanco-Vásquez FE, García-Arroyave LP. [General Directorate of Research, University of San Carlos of Guatemala: Reproductive Aspects of *Cynoscion reticulatus* and *Micropogonias altipinnis* from Artisanal Fishing Landings in Sipacate, Pacific of Guatemala]. 2021 Feb; Report No.: 2020-23.
21. Diario Oficial de la Federación. [Agreement Approving the Update of the National Fisheries Charter and Its Annex]. Secretaría de Agricultura, Ganadería, Desarrollo Rural, Pesca y Alimentación. (2004 Mar 15). 2004. Available from: https://dof.gob.mx/nota_detalle.php?codigo=680061&fecha=15/03/2004#gsc.tab=0
22. Muro-Torres VM, Soto-Jiménez MF, Amezcua F, Green L, Quintero J. Food web structure of a subtropical coastal lagoon. Aquat Ecol. 2019;53:407-430. <https://doi.org/10.1007/s10452-019-09698-0>
23. Arreguín-Sánchez F, Arcos E, Chávez EA. Flows of biomass and structure in an exploited benthic ecosystem in the gulf of California, Mexico. Ecol Modell. 2002;156:67-183. [https://doi.org/10.1016/S0304-3800\(02\)00159-X](https://doi.org/10.1016/S0304-3800(02)00159-X)
24. García-Rodríguez FJ, Aurióles-Gamboa D. Spatial and temporal variation in the diet of the California sea lion (*Zalophus californianus*) in the Gulf of California, Mexico. Fish Bull. 2004;102:47-62.
25. Mellink E, Romero-Saavedra AL. Diet of California sea lions, *Zalophus californianus*, at San Jorge Island, northern Gulf of California, Mexico, 1998–1999. Cienc Mar. 2005;31(2):369-377.
26. Fricke R, Eschmeyer NW, Fong JD. [Internet]. Eschmeyer's Catalog of Fishes. [cited 2022 Aug 16]. Available from: <https://researcharchive.calacademy.org/research/ichthyology/catalog/SpeciesByFamily.asp>
27. Zar JH. Biostatistical analysis. New Jersey; Pearson Prentice Hall. 2010.
28. Rojas-Bracho L, Reeves RR, Jaramillo-Legorreta A. Conservation of the vaquita *Phocoena sinus*. Mammal Review. 2006;36(3):179-216.
29. Parenti P, Randall JE. An annotated checklist of the fishes of the family Serranidae of the world with description of two new related families of fishes. Fishtaxa-Journal of Fish Taxonomy. 2020; (15).
30. Summers AP. Stiffening the stingray skeleton—an investigation of durophagy in myliobatid stingrays (Chondrichthyes, Batoidea, Myliobatidae). J Morphol. 2000;243(2):113-126. [https://doi.org/10.1002/\(SICI\)1097-4687\(200002\)243:2%3C113::AID-JMOR1%3E3.0.CO;2-A](https://doi.org/10.1002/(SICI)1097-4687(200002)243:2%3C113::AID-JMOR1%3E3.0.CO;2-A)
31. Oufiero CE, Holzman RA, Young FA, Wainwright PC. New insights from serranid fishes on the role of trade-offs in suction-feeding diversification. J Exp Biol. 2012;215(21):3845-3855. <https://doi.org/10.1242/jeb.074849>

32. Ramírez-Díaz C, Peña R, Diogo R, Cruz-Escalona VH. Comparative cranio-mandibular myology of three species of Batoidea from the Southern Gulf of California, Mexico. *J Morphol*. 2023;284(1):e21547.
33. Wilga CD, Motta PJ, Sanford CP. Evolution and ecology of feeding in elasmobranchs. *Integr Comp Biol*. 2007;47(1):55-69. <https://doi.org/10.1093/icb/icm029>
34. Barros NB, Wells RS. Prey and Feeding Patterns of Resident Bottlenose Dolphins (*Tursiops truncatus*) in Sarasota Bay, Florida, J. Mammal. 1998; 79(3):1045–1059. <https://doi.org/10.2307/1383114>
35. De Roy T, Espinoza ER, Trillmich F. Cooperation and opportunism in Galapagos sea lion hunting for shoaling fish. *Ecol Evol*. 2021. <https://doi.org/10.1002/ece3.7807>
36. Hansen MJ, Kurvers RHJM, Licht M, Häge J, Pacher K, Dhellemmes F, et al. California sea lions interfere with striped marlin hunting behaviour in multi-species predator aggregations. *Philos Trans R Soc Lond B Biol Sci*. 2023;378(1878):20220103. <https://doi.org/10.1098/rstb.2022.0103>
37. Álvarez-León R. A specimen of *Lutjanus argentiventris* (Peters) lacking pelvic fins, México. *Anales Inst Biol Univ Nac Autón México*. 1983;54:229-35. Available from: https://www.researchgate.net/publication/230325483_A_specimen_of_Lutjanus_argentiventris_Peters_lacking_pelvic_fins
38. Mariasingarayan Y, Danaraj J, Veeraiyan B, Fjellidal PG, Karuppiyah K, Narayanasamy R. Vertebral column deformity in six species of wild fish at the Coromandel coast, Bay of Bengal India. *Acuac Fish*. 2022. <https://doi.org/10.1016/j.aaf.2022.05.004>
39. Sindermann CJ. Pollution-associated diseases and abnormalities of fish and shellfish: a review. *Fish Bull*. 1979;76:717-749. [cited 2023 Feb 16]. Available from: <https://fisherybulletin.nmfs.noaa.gov/sites/default/files/pdf-content/fish-bull/sindermann%20%281%29.pdf>
40. Walker W. (1981). Geographical variation in morphology and biology of bottlenose dolphins (*Tursiops*) in the Eastern North Pacific. NOAA/NMFS Southwest Fisheries Center Administrative; Jan 1981. Report. No. LJ-81-03C. Contract No. 03-7-208-35238.
41. Cockcroft, V, Ross G. Food and feeding of the Indian Ocean bottlenose dolphin off Southern Natal, South Africa. In: Leatherwood S, Reeves RR, editors. *The bottlenose dolphin*. California: Academic Press. 1990; p. 295-308.
42. Barros N, Odell D. Food habits of bottlenose dolphins in the south-eastern United States. In: Leatherwood S, Reeves RR, editors. *The bottlenose dolphin*. California: Academic Press. 1990; p. 309-328.
43. Waerebeek K, Reyes J, Read A, McKinnon J. Preliminary observations of bottlenose dolphins from the Pacific Coast of South America. In: Leatherwood S, Reeves RR, editors. *The bottlenose dolphin*. California: Academic Press. 1990; 1990; p. 143-154.

44. Ronje EI, Barry KP, Sinclair C, Grace MA, Barros N, Allen J, Balmer B, Panike A, Toms C, Mullin KD, Wells RS. A common bottlenose dolphin (*Tursiops truncatus*) prey handling technique for marine catfish (Ariidae) in the northern Gulf of Mexico. PloS one. 2017;12(7). <https://doi.org/10.1371/journal.pone.0181179>



© 2024 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY NC) license (<https://creativecommons.org/licenses/by-nc/4.0/>).